

Dual-species origin of an adaptive chemical defense polymorphism

Kenneth M. Olsen¹ , David M. Goad^{1*} , Sara J. Wright^{1,2*}, Maya L. Dutta¹, Samantha R. Myers¹, Linda L. Small¹ and Lin-Feng Li³

¹Department of Biology, Washington University in St. Louis, 1 Brookings Dr., St Louis, MO 63130, USA; ²Biological Sciences Department, Rowan University, Glassboro, NJ 08028, USA;

³Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, School of Life Sciences, Fudan University, Shanghai 200438, China

Author for correspondence:

Kenneth M. Olsen

Email: kolsen@wustl.edu

Received: 7 April 2021

Accepted: 22 July 2021

New Phytologist (2021)

doi: 10.1111/nph.17654

Key words: allopolyploid speciation, chemical defense, cyanogenic glucoside, intraspecific polymorphism, linamarase, *Trifolium repens* (Fabaceae).

Summary

- Allopolyploid speciation and chemical defense diversification are two of the most characteristic features of plant evolution; although the former has likely shaped the latter, this has rarely been documented. Here we document allopolyploidy-mediated chemical defense evolution in the origin of cyanogenesis (HCN release upon tissue damage) in white clover (*Trifolium repens*).
- We combined linkage mapping of the loci that control cyanogenesis (*Ac*, controlling production of cyanogenic glucosides; and *Li*, controlling production of their hydrolyzing enzyme linamarase) with genome sequence comparisons between white clover, a recently evolved allotetraploid, and its diploid progenitors (*Trifolium pallescens*, *Trifolium occidentale*).
- The *Ac* locus (a three-gene cluster comprising the cyanogenic glucoside pathway) is derived from *T. occidentale*; it maps to linkage group 2O (*occidentale* subgenome) and is orthologous to a highly similar cluster in the *T. occidentale* reference genome. By contrast, *Li* maps to linkage group 4P (*pallescens* subgenome), indicating an origin in the other progenitor species.
- These results indicate that cyanogenesis evolved in white clover as a product of the interspecific hybridization that created the species. This allopolyploidization-derived chemical defense, together with subsequent selection on intraspecific cyanogenesis variation, appears to have contributed to white clover's ecological success as a globally distributed weed species.

Introduction

A hallmark of plant species diversification has been the evolution of diverse specialized metabolites, many of which function as antiherbivore chemical defenses (Pichersky & Lewinsohn, 2011; Mithöfer & Boland, 2012). In recent decades, advances in genomics, combined with biochemical and physiological studies, have provided important insights into the molecular bases of these defense metabolite pathways and their macroevolutionary origins (e.g. Zhuang *et al.*, 2012; Edger *et al.*, 2015; Xu *et al.*, 2017; Lai *et al.*, 2020). On a microevolutionary timescale, species that are polymorphic for chemical defense production have served as especially valuable systems for studying the molecular basis of adaptive variation and the roles of natural selection and other forces in maintaining it (Brachi *et al.*, 2015; Wager & Li, 2018; Keith & Mitchell-Olds, 2019; Lowry *et al.*, 2019).

Like chemical defense evolution, another defining feature of plant diversification has been allopolyploid speciation, whereby interspecific hybridization and retention of both parental species' genomes creates a new polyploid daughter species. This allopolyploidization process has the potential to generate novel adaptive

variation for diverse traits including chemical defense metabolites (Soltis *et al.*, 2014; Van de Peer *et al.*, 2017). Experiments with synthetically created allopolyploids have demonstrated, for example, that distinct chemical defense profiles can arise through remodeling of parental species' gene expression networks and post-transcriptional processing (Anssour & Baldwin, 2010). However, studies are largely lacking on the role of allopolyploidization for chemical defense evolution in natural wild species; in particular, there is little empirical work on the relationship between chemical defense traits in allopolyploids and the genetic contributions of their diploid progenitors. Given the likely role of defense metabolite evolution in angiosperm diversification (Pichersky & Lewinsohn, 2011; Mithöfer & Boland, 2012), and the prevalence of allopolyploid speciation in plants (Van de Peer *et al.*, 2017), characterizing this relationship is an important step towards understanding the mechanisms of adaptive diversification in the plant kingdom.

A long-standing model system for plant chemical defense evolution is cyanogenesis (HCN release following tissue damage) and the polymorphism for this trait that occurs in white clover (*Trifolium repens* L.). First described more than a century ago (Armstrong *et al.*, 1913), white clover plants vary in the cyanogenic response, with both cyanogenic and acyanogenic

*These authors contributed equally to this work.

genotypes found in wild populations. Latitudinal and elevational clines in cyanogenesis have evolved in white clover throughout the native species range in Europe and in introduced populations worldwide; cyanogenic plants predominate in warmer and drier locations, but are largely absent from populations at higher latitudes and elevations (Daday, 1954, 1958; Kooyers & Olsen, 2012, 2013). Since the 1950s, numerous studies have attempted to identify the selective factors that maintain this polymorphism and favor the recurrent evolution of climate-associated cyanogenesis clines. Key factors likely include herbivore deterrence, energetic costs of producing the defense and abiotic stress adaptation (reviewed by Hughes, 1991; Olsen *et al.*, 2013; Kooyers *et al.*, 2018). More recently, the molecular basis of the white clover cyanogenesis polymorphism also has been characterized (Olsen *et al.*, 2007, 2008; Olsen & Small, 2018). This chemical defense polymorphism has come to serve as a textbook example of intraspecific variation maintained by geographically heterogeneous selection (Briggs & Walters, 2016; Futuyama & Kirkpatrick, 2017), and recently it also has been adopted as a model for understanding adaptation to urban environments (Thompson *et al.*, 2016; Santangelo *et al.*, 2020). White clover is a common weed species worldwide; the fact that climate-associated cyanogenesis clines have evolved repeatedly throughout the native and introduced species range suggests that selection on this chemical defense polymorphism has facilitated its global expansion and widespread ecological success (Daday, 1958; Kooyers & Olsen, 2013; Santangelo *et al.*, 2020).

White clover is an allotetraploid species ($2n=4x=32$), having arisen through the hybridization of two diploid ($2n=2x=16$) *Trifolium* species (Williams *et al.*, 2012; Griffiths *et al.*, 2019), and a critical unanswered question for the cyanogenesis polymorphism is how the evolution of this chemical defense relates to the species' hybrid origin. The cyanogenic response occurs through the interaction of two chemical precursors that are separated in intact tissue and brought together upon tissue damage: cyanogenic glucosides (CNGlcs), which in leaf tissue are likely compartmentalized in the vacuole, and their hydrolyzing enzyme linamarase, which is present in the apoplast (Kakes, 1985; Gleadow & Møller, 2014). Inheritance of the polymorphism is determined by two unlinked simple Mendelian polymorphisms that control the production of each precursor: *Ac/ac* controls the presence/absence of CNGlcs, and the independently segregating *Li/li* gene controls the presence/absence of linamarase (Hughes, 1991). For both polymorphisms, homozygote recessive genotypes (*acac* or *lili*) lack the corresponding chemical precursor. Thus, plants that carry at least one dominant allele at both genes have both required precursors and are cyanogenic (referred to as the AcLi 'cyanotype'), whereas plants that are homozygous recessive at either or both genes are acyanogenic (acLi, Acli and acli 'cyanotypes').

At the molecular level, the *Ac/ac* polymorphism corresponds to a genomic presence/absence polymorphism for the three-gene metabolic cluster that constitutes the CNGlc biosynthesis pathway (Olsen & Small, 2018). In parallel, the unlinked *Li/li* polymorphism corresponds to a genomic presence/absence polymorphism for the gene encoding the linamarase protein (Olsen *et al.*, 2007). For both *Ac/ac* and *Li/li*, the recessive

(gene-deletion) alleles have arisen multiple times independently in white clover (Olsen *et al.*, 2013), and gene deletions at these same loci have evolved in parallel in several related *Trifolium* species (Olsen *et al.*, 2014; Olsen & Small, 2018). Because inheritance of both the *Ac/ac* and *Li/li* polymorphisms follows a single gene, two-allele model, both of the cyanogenesis loci have been presumed to be present within only one of white clover's two parental subgenomes (Kakes & Hakvoort, 1994; Badr *et al.*, 2002). However, their genomic locations have remained unresolved; this has left open the questions of whether both genes are derived from the same diploid progenitor, and which progenitor(s) contributed them.

The recent publication of a white clover reference genome, together with conclusive identification of its diploid progenitor species (Williams *et al.*, 2012; Griffiths *et al.*, 2019), has opened up the possibility of tracing the evolutionary origins of the cyanogenesis loci. *T. repens* is estimated to have originated in the very recent evolutionary past (~15–28 000 yr ago) by multiple hybridizations between *Trifolium palleescens*, a species now distributed in alpine regions of central Europe, and *Trifolium occidentale*, a species currently restricted to Atlantic coastal margins of Western Europe (Griffiths *et al.*, 2019). Curiously, the present-day occurrence of cyanogenesis in these diploid progenitors provides only limited insight for inferring the trait's origins. *T. occidentale* carries the *Ac/ac* polymorphism (Kakes & Chardonnens, 2000), and, as in white clover, *ac* alleles correspond to deletions of the *Ac* gene cluster (Olsen *et al.*, 2014; Olsen & Small, 2018). However, extensive surveys of wild populations have not detected linamarase (or the *Li* gene) in this species (Kakes & Chardonnens, 2000). For *T. palleescens*, neither of the cyanogenic precursors (or their underlying genes) have been detected in population sampling to date (Gibson *et al.*, 1972; Olsen *et al.*, 2014) (see also Supporting Information Table S1).

Because gene sequences of the two diploid progenitor species are highly similar to each other ($\geq 97\%$ identity on average; Table S2) and also are highly conserved with their respective descendant subgenomes in *T. repens* (Griffiths *et al.*, 2019), most white clover sequence contigs cannot be definitively assigned to one subgenome or the other in the currently available shotgun assembly reference genome (A. Griffiths and R. Moraga, pers. comm.; D. M. Goad, unpublished). Thus, the *Ac* and *Li* gene sequences alone are insufficient to determine their subgenome locations and diploid contributors. This problem can potentially be overcome, however, by combining sequence analyses with genetic mapping of the *Ac/ac* and *Li/li* polymorphisms and integration of linkage map data into the genome assembly. The availability of draft genome assemblies for each diploid progenitor species (Griffiths *et al.*, 2019) further facilitates this genetic mapping process.

In this study we generated high-density linkage maps from two white clover F_2 mapping populations and combined genetic mapping of the *Ac/ac* and *Li/li* cyanogenesis polymorphisms with gene and genome sequence analyses to assess the evolutionary origins of the cyanogenesis loci. We addressed the following questions: (1) Given that the *Ac* gene cluster is known to exist in *T. occidentale*, do genetic map data and genome sequence analyses

support the hypothesis that the white clover *Ac* locus is derived from that diploid species? (2) Is the *Li* locus located in the same subgenome as *Ac*, indicating that both cyanogenic components were contributed by the same diploid progenitor? Our results reveal that this textbook example of a plant chemical defense has a dual-species origin, with the required cyanogenesis genes contributed through the interspecific hybridization that created this widely adapted allopolyploid species.

Materials and Methods

Generation of mapping populations

Trifolium repens is a native species of Europe that has been introduced into mesic, temperate regions worldwide; it is grown for forage and is widely naturalized in lawns, pastures and other disturbed habitats. For this study, two *T. repens* F₂ mapping populations were generated from biparental crosses using three wild-collected North American plants. The parental genotypes were selected such that the presence/absence of cyanogenic glucosides (CNGlcs) (*Ac/ac*) and of linamarase (*Li/li*) polymorphisms would segregate to create all four cyanotypes in the F₂ generation: DMN_010 (*acac, lili*; originating from Duluth, MN); STL_0701 (*acac, lili*; from St Louis, MO) (Kooyers & Olsen, 2012); and GFL_007 (*AcAc, LiLi*; from Gainesville, FL). GFL_007 served as a parent in both mapping populations, yielding the DMN_010 × GFL_007 (hereafter DG) and STL_0701 × GFL_007 (hereafter SG) populations. For both populations, hand crosses were performed in the glasshouse between parents in both directions to generate 50–100 F₁ genotypes; the species has gametophytic self-incompatibility (Casey *et al.*, 2010), so no self-fertilization occurred. Within each F₁ population, random cross-pollinations were performed to generate *c.* 500 F₂ genotypes per population (Tables S3, S4). For the DG population, solitary bees (Crown Bees, Woodinville, WA) were used to cross-pollinate F₁ plants in enclosed cages in the greenhouse. For the SG population, random crosses were performed by hand to produce F₂ seed.

The three parental genotypes, along with F₁ and F₂ genotypes, were grown in the Washington University glasshouse facilities under standard conditions (16 h : 8 h, light : dark photoperiod with 140 μmol supplemental lighting as needed; 18–23°C; *c.* 50% relative humidity). Seeds were scarified using fine grit sandpaper and placed on mist benches in the greenhouse for germination. Seedlings were transferred to individual pots for establishment after producing at least one true leaf.

Genotyping

For all plants of all generations in both mapping populations, young fresh leaf tissue (120–150 mg) was ground to powder in liquid N₂ using a mortar and pestle, and high-purity genomic DNA (gDNA) was extracted using a protocol modified from Whitlock *et al.* (2008). DNA was run on 0.8% agarose gels to check for quality and integrity. Samples were quantified and standardized to 17–23 ng μl⁻¹ as measured by a Qubit[®] 2.0 Fluorometer (Invitrogen).

***Ac* and *Li* polymorphisms** Parental, F₁ and F₂ plants were genotyped for the *Ac/ac* and *Li/li* gene presence/absence polymorphisms using previously described PCR genotyping protocols (Olsen *et al.*, 2007, 2008) (Table S5). Negative PCR results were repeated at least once to confirm recessive (gene-absence) genotypes. Genotypes were further confirmed by cyanogenesis phenotyping using leaf tissue and Feigl–Anger HCN tests (Feigl & Anger, 1966), as described previously (Olsen *et al.*, 2007). For acyanogenic genotypes, the presence of individual cyanogenic components was assessed by exogenously adding complementary reagents (either linamarase or cyanogenic glucosides) individually to leaf tissue and re-testing with Feigl–Anger paper following the previously described protocol. All F₁ genotypes in both populations were confirmed to be cyanogenic (AcLi cyanotype), consistent with the expectation that they would be heterozygous at both cyanogenesis genes (*Acac, Lili*).

Genotyping-by-sequencing Standardized quantities of gDNA samples from the parent plants and all F₁ and F₂ plants were prepared in randomized libraries, each containing 95 samples and one negative control sample (water), using a genotyping-by-sequencing (GBS) protocol modified from (Huang *et al.*, 2014) with 100 ng gDNA and *ApeKI* for digestion (Elshire *et al.*, 2011). Common and adapter barcode sequences were created using an online GBS barcode generator (Deena Bioinformatics, Thomas van Gorp, Nijmegen, the Netherlands) for design of oligonucleotide barcode sequences (IDT, Coralville, IA, USA) (Table S6). Following enzyme digestion and adapter ligations, cleanup and size-selection steps were performed using Agencourt AMPure XP beads (Beckman Coulter, Brea, CA, USA). Samples then were pooled and PCR-amplified using primers from Elshire *et al.* (2011) (Table S5).

Paired-end sequencing (150-bp reads) was performed with each library in a unique lane using the Illumina Hi-Seq 2500 platform (Novogene Corp., Chula Vista, CA, USA). GBS libraries (13 in total) were run individually on sequencing lanes with a 15% PhiX Control spike-in (Illumina, San Diego, CA, USA). The three parents of the mapping populations were included as standards in all 13 libraries to increase read depth and reliability of heterozygous single nucleotide polymorphism (SNP) calls for optimal selection of parental markers for genetic mapping.

Raw GBS reads were analyzed using an in-house version of the FAST-GBS pipeline (Torkamaneh *et al.*, 2017) that was modified to optimize handling of paired-end data. Briefly, reads were demultiplexed with SABRE (<https://github.com/najoshi/sabre>) and trimmed with CUTADAPT (Martin, 2011). Reads then were mapped back to the white clover reference genome (Griffiths *et al.*, 2019) using BWA (Li & Durbin, 2009) on default paired-end settings. Parental and F₁ SNP variants were called with *Platypus* (Rimmer *et al.*, 2014), which filtered reads with unmapped or distant mates.

Linkage map construction

In order to generate a set of markers that were most likely to segregate in a 1 : 2 : 1 ratio in the F₂ populations, SNPs were filtered

to the subset that were homozygous in both parents and had >0.7 heterozygosity in the F₁ population. Allele counts for each F₂ SNP were passed to GUSMAP v.1.0 (Bilton *et al.*, 2018), where additional SNP filtering was performed before linkage map construction. Any F₂ lines with <15× average read depth were excluded (Tables S2, S3). Filtering also removed any markers that did not meet the following criteria: minor allele frequency (MAF) >0.05, missing data <0.1, average read depth >5×, or a *P*-value <0.01 on a genotype frequency test (indicating deviations from 1:2:1 segregation in the F₂ populations). Additionally, SNPs were pruned such that only one SNP was present from any given 2000-bp region to represent a single GBS fragment.

The *createLG* function in GUSMAP was used to calculate pairwise *r*² and logarithm of odds (LOD) scores; SNPs were assigned to a linkage group if their LOD score in that group exceeded 45. Because GUSMAP v.1.0 was designed for data from an F₁ pseudotestcross population, this function was modified slightly to use only SNPs that were informative in both parents rather than those that are only maternally or paternally informative. The SNPs then were ordered using the *orderLG* function with default parameters. Markers that appeared to be incorrectly placed upon manual inspection of recombination rates were removed, and linkage maps were created using the *computeMap* function with the Kosambi method for calculating distances. Markers that introduced large gaps were then removed from the map and distances were recalculated.

In order to assign labels to linkage groups that were consistent with previously published white clover linkage maps (Barrett *et al.*, 2004; Griffiths *et al.*, 2013), we generated a 200-bp sequence for each marker based on the reference genome sequence 100 bp upstream and downstream of each SNP; these sequences were then used as BLAST queries against the reference genomes of the two progenitor species and *Medicago truncatula*. Linkage groups were assigned to either the O or P subgenome based on which progenitor species genome the most markers from a given linkage group matched with highest-scoring BLAST hits (Table S7). Within a subgenome, linkage groups were labeled to match previous published linkage maps based on shared synteny to *M. truncatula* chromosomes (Fig. S1) (Griffiths *et al.*, 2013). The marker order for a given linkage group was inverted where necessary to match its direction in the previously published maps.

Genetic mapping

The *Ac/ac* and *Li/li* polymorphisms were mapped separately in each F₂ population to identify the locations of the corresponding loci in the DG and SG linkage maps. F₂ genotypes were not used in the genetic mapping analysis if they were missing >75% genotype data; this left 423 and 500 genotypes in the DG and SG populations, respectively (Tables S3, S4). Mapping analysis was performed with the R/QTL package (Broman *et al.*, 2003). The *scanone* function was used to map quantitative trait loci (QTLs) for *Ac/ac*, *Li/li* and cyanogenesis. QTLs were considered significant if their LOD score was higher than a *P*=0.05 confidence threshold, generated for each phenotype from 1000 permutations

of the data. For each QTL, LOD scores and the percentage variance explained (PVE) were calculated with the *fitqtl* function, and 1.5-LOD drop intervals were calculated with the *lodint* function. For cyanogenesis, which had two significant peaks, we fit a two-QTL model with refined positions from the *refineqtl* function. All tests were performed with a binary trait model and the Haley–Knott algorithm (Haley & Knott, 1992).

Cyanogenesis gene sequence analysis

The previously published 138-kb *Ac* gene sequence (GenBank accession MH059954, comprising the three-gene CNg1c metabolic cluster) was used in BLAST analyses on the NCBI platform, querying whole genome shotgun contigs (WGS database) of each diploid progenitor species to identify sequences with highest identity to the white clover sequence. Alignment to the best hit was visualized using the COGE web platform (<https://genomevolution.org/coge/GEvo.pl>) with default parameter settings. Likewise, the 3.9-kb *Li* gene sequence was used as a query in BLAST analyses of the *T. occidentale* and *T. pallescens* genomes. To assess potential similarity in the genomic sequence flanking the *Li* locus to progenitor genome sequences, BLAST analyses were also performed using a larger 77.6-kb genomic sequence containing the *Li* gene that was derived from a white clover BAC library (Olsen & Small, 2018). To determine whether the *Ac* and *Li* loci are present in the *T. repens* shotgun genome assembly, both loci also were used as queries in BLAST analysis against that reference genome in the NCBI portal.

Results

White clover linkage maps

Full-sib crosses among F₁ plants of each parental cross (DMN_010 × GFL_007 and STL_0701 × GFL_007) yielded 502 and 500 F₂ individuals for the DG and SG populations, respectively. After filtering for the subset of SNPs that could be mapped with paired-end sequence reads and that were homozygous in both parents of a mapping population but heterozygous in F₁ plants (see Materials and Methods), the DG mapping population was characterized by 8095 polymorphic SNP markers and the SG population was characterized by 7665 polymorphic SNPs. Filtering out of any F₂ genotypes with <15× coverage yielded 252 lines in the DG population and 191 lines in the SG population that were used in linkage map construction (Tables S3, S4). For both datasets, linkage maps were composed of 16 linkage groups, consistent with white clover's haploid chromosome number (*n*=2*x*=16), with eight linkage groups corresponding to the *T. occidentale* subgenome (chromosomes 1O to 8O, following the nomenclature of Griffiths *et al.*, 2019) and the other eight corresponding to the *T. pallescens* subgenome (chromosomes 1P to 8P) (Fig. 1a,b).

The linkage map for the DG F₂ population included 2575 SNPs with a total length of 5057.6 cM and average marker interval of 2.0 cM (Table S8). Of these markers, 1104 mapped to the *T. occidentale* subgenome and 1471 mapped to the *T. pallescens*




Fig. 1 Linkage maps and cyanogenesis quantitative trait loci (QTL) locations in the DG (a) and SG (b) mapping populations in white clover (*Trifolium repens*). Linkage groups are identified by subgenome chromosome number, where O and P designations correspond (respectively) to the *Trifolium occidentale* and *Trifolium pallescens* subgenomes. Horizontal lines indicate genetic map locations of genotyping-by-sequencing markers. Colored bars indicate locations and approximate span of significant QTLs for cyanogenic glucoside (CNglc) production (*Ac*), linamarase production (*Li*) and cyanogenesis.

subgenome. The linkage map for the SG F₂ population included 2437 SNPs with a total length of 5815.6 cM and average interval of 2.4 cM (Table S9). Of these markers, 1045 were mapped to chromosomes belonging to the *T. occidentale* subgenome and 1392 mapped to the *T. pallescens* subgenome. The SNP markers in both mapping populations thus provide roughly equal representation of white clover's two parental subgenomes.

QTL mapping of cyanogenesis loci

In both mapping populations, the *Ac* gene mapped with high confidence to a single, highly localized genomic region on the top half of chromosome 2O (LOD = 85.2 and PVE = 54.2% in DG; LOD = 120.7, PVE = 67.1% in SG) (Figs 1a,b, S2, S3; Table 1). Likewise, *Li* mapped to a single, highly localized region on the bottom of chromosome 4P in both populations, again with high confidence (LOD = 90.8 and PVE = 56.5% in DG;

LOD = 103.2, PVE = 61.3% in SG) (Figs S4, S5). The identification of a single QTL peak for each cyanogenesis gene is consistent with the well-established simple Mendelian inheritance of the *Ac/lac* and *Li/li* polymorphisms. Correspondingly, the cyanogenesis phenotype mapped to the locations of the *Ac* and *Li* loci, and only to those two locations, with high confidence in both populations (> 70 LOD) (Figs S6, S7). Notably, the location of the *Ac* locus within the O subgenome and the *Li* locus within the P subgenome strongly suggests that white clover's two cyanogenesis components were inherited from different diploid progenitors, with CNglc production derived from *T. occidentale* and linamarase production inherited from *T. pallescens*.

Genome sequence analyses

***Ac* locus** Sequence comparisons of the white clover *Ac* locus to shotgun genome assemblies of the two progenitor species were

Table 1 Quantitative trait loci (QTL) mapping results for cyanogenic glucoside production (*Ac*), linamarase production (*Li*) and cyanogenesis phenotypes for the DG and SG mapping populations in white clover (*Trifolium repens*).

Population	Phenotype	Linkage group	Position (cM)	LOD score	PVE	1.5-LOD drop interval			
						Start (cM)	Stop (cM)	First marker	Last marker
DG	<i>Ac</i> (CNglcs)	2O	119.9	85.2	54.24%	118.07	120.23	DG_2_061	DG_2_063
DG	<i>Li</i> (linamarase)	4P	376.9	90.8	56.51%	375.16	378.12	DG_12_151	DG_12_163
DG	Cyanogenesis	2O	119.9	66.61	31.52%	116.66	120.23	DG_2_060	DG_2_063
DG	Cyanogenesis	4P	376.9	70.38	33.94%	374.03	378.12	DG_12_148	DG_12_163
SG	<i>Ac</i> (CNglcs)	2O	79.73	120.71	67.10%	74.73	84.89	SG_2_033	SG_2_036
SG	<i>Li</i> (linamarase)	4P	318.1	103.2	61.34%	313.00	318.93	SG_12_142	SG_12_144
SG	Cyanogenesis	2O	83.8	95.72	40.84%	74.73	84.89	SG_2_033	SG_2_036
SG	Cyanogenesis	4P	318.1	73.71	28.05%	313.00	318.93	SG_12_142	SG_12_144

cM, centimorgan; CNglcs, cyanogenic glucosides; LOD, logarithm of odds; PVE, percentage variation explained.

consistent with genetic mapping results in suggesting inheritance from *T. occidentale*. BLAST analysis revealed that much of the 138-kb *Ac* gene cluster, including both coding and noncoding sequence, was nearly identical to a single 169.2-kb region within the *T. occidentale* genome ($\geq 97\%$ sequence identity across 78.2% of the region) (Fig. 2; Table S10). All three genes of the CNglc biosynthetic pathway fell within highly conserved regions (97.5%, 99.2% and 99.6% sequence identity for *CYP79D15*, *CYP736A187* and *UGT85K17* respectively), and synteny also was preserved across the entire gene cluster. No significant matches were detected for any of the CNglc synthesis genes outside of this 169.2 kb sequence in the *T. occidentale* genome assembly. These results suggest that the white clover *Ac* locus is likely orthologous to, and thus descended from, the highly similar gene cluster in the *T. occidentale* genome.

By contrast with the high similarity detected in *T. occidentale*, the white clover *Ac* sequence showed only negligible similarity to *T. pallescens* genomic sequences. Across the entire 138 kb *Ac* region, the longest *T. pallescens* sequence with $\geq 97\%$ identity was a 692-bp fragment that corresponds to a previously described *Ty1/copia*-like retrotransposon that occurs within the gene cluster (Olsen & Small, 2018) (Table S10). Among the three CNglc synthesis genes, the closest match was for a 173-bp portion of the 2.0-kb *CYP736A187* gene which showed 82.1% identity to a *T. pallescens* sequence. Taken together with the genetic mapping data and the *T. occidentale* sequence comparisons, these results strongly suggest that the white clover *Ac* locus is derived from *T. occidentale* and not from *T. pallescens*.

Li locus Unlike inferences for *Ac*, genome sequence analyses were inconclusive in identifying the ancestral sequence of the white clover *Li* gene. BLAST analysis of the 3.8-kb *Li* gene against the *T. occidentale* genome yielded a single best hit with 92.1% sequence identity. However, this *T. occidentale* sequence shows greater similarity (97.1% identity) to a previously described *Li*-paralog in the white clover genome that does not function in cyanogenesis (Olsen *et al.*, 2007) (Table S10); the gene is, thus, more likely to be the *T. occidentale* ortholog of the non-cyanogenic *Li*-paralog than the *Li* gene. Likewise, BLAST queries of the *T. pallescens* genome with *Li* revealed a single best hit; as

with *T. occidentale*, the sequence showed higher nucleotide similarity to the white clover noncyanogenic *Li*-paralog than to *Li* (96.0% vs 94.6% identity, respectively; Table S10). BLAST searches using a larger 77.6 kb sequence containing the *Li* gene also did not yield any obvious close matches within either diploid genome, with best hits for >1-kb fragments showing $\leq 86\%$ identity to both species. Thus, although genetic mapping of the white clover *Lilli* polymorphism indicates its placement in the *T. pallescens* subgenome, the *Li* gene sequence does not appear to be present in the published reference genome of that diploid progenitor species. This result suggests that present-day *T. pallescens* either lacks the *Li* gene altogether or that the species is polymorphic for *Lilli*, with the published genome derived from a *lili* (gene-deletion homozygote) genotype. Consistent with both of these possibilities, cyanogenesis genotyping and phenotyping of the *T. pallescens* accession used to create the reference genome (AZ 1895) indicated that the *Li* gene is not present in that genotype (Table S1).

Trifolium repens reference genome comparisons BLAST analysis of the *Ac* locus against the *T. repens* reference genome confirmed the presence of the gene cluster in the accession used for genome sequencing (100% identity for a 139.1-kb region; Table S10). The *T. repens* contig containing the *Ac* locus currently is assigned to *T. repens* chromosome 1P; given the lack of congruence with genetic map data, where *Ac* is placed unambiguously at the top of 2O (Fig. 1; Table 1), this designation likely is an artifact of contig misassignment in the current shotgun-assembly draft genome. The *Li* locus also was confirmed to be present in the white clover reference genome (100% identity for the 3.8-kb gene; Table S10). In this case, the *Li*-containing region appears to be assigned to the correct chromosome in the draft assembly; it is annotated as a chromosome 4P sequence, consistent with QTL mapping results (Fig. 1; Table 1).

Discussion

Two of the most characteristic features of plant species diversification are speciation via allopolyploidy and the evolution of diverse chemical defense metabolites. Here we have examined




Fig. 2 Alignment of the white clover (*Trifolium repens*) *Ac* locus for cyanogenic glucoside (CNgIc) production (upper) against the *Trifolium occidentale* shotgun genome assembly (lower). Shaded connectors indicate regions of $\geq 97\%$ sequence identity. Colored regions correspond to the three genes of the cyanogenic glucoside biosynthetic pathway; arrows indicate transcript orientation. Crossed-hatched boxes (*T. occidentale*) indicate regions of undetermined sequence in the draft genome. The *T. occidentale* region corresponds to GenBank accession CM022957, sequence range: 59 820 332–59 989 571.

how the former phenomenon gave rise to a textbook example of the latter, namely cyanogenesis in the allotetraploid species white clover (*T. repens*). Through the integration of linkage maps and genetic mapping of the white clover presence/absence of cyanogenic glucosides (CNgIcs) (*Ac/ac*) and of linamarase (*Lilli*) polymorphisms with genome sequence analyses in its two diploid progenitor species, we have documented a dual-species origin for the chemical defense in this allotetraploid species. Specifically, the synthesis of CNgIcs, which requires the three-gene metabolic gene cluster comprising the *Ac* locus, has been contributed by *T. occidentale* (Fig. 1a,b; Table 1); sequence comparisons between *T. repens* and *T. occidentale* reveal a high degree of conservation across the entire gene cluster (Fig. 2). For the hydrolyzing enzyme, linamarase, which is required to liberate HCN from CNgIcs, genetic mapping indicates that the underlying *Li* gene has an origin in the *T. pallescens* genome, although the gene is absent in the genotype used to generate the published reference genome (Table S1). Our confidence in these evolutionary inferences is bolstered by the fact that white clover's two subgenomes show very high genome integrity with their respective diploid progenitor species, with $\leq 5\%$ gene loss per subgenome and very limited interhomeologous recombination since polyploidization (Griffiths *et al.*, 2019); this high degree of subgenome conservation suggests that the linkage map data and genome sequence comparisons are both accurately reflecting the cyanogenesis genes' evolutionary origins. Support for these conclusions is further strengthened by the very high degree of similarity between our *Trifolium–Medicago* synteny plots (Fig. S1) and those reported in previous analyses (e.g. Griffiths *et al.*, 2019). Below we discuss these findings in the context of the white clover cyanogenesis polymorphism, *Trifolium* evolution, and allopolyploid adaptation more broadly.

Cyanogenic glucosides and the *Ac* locus

As a chemical defense that requires physical interaction between two normally separated biochemical precursors, cyanogenesis has been described as a 'cyanide bomb' (Morant *et al.*, 2008) that is triggered by chewing herbivores. By this analogy, the 'explosives' component of the white clover cyanide bomb (i.e. CNgIcs) was contributed by the diploid progenitor *T. occidentale*. This coastal

western European species is not cyanogenic but does carry the *Ac/ac* polymorphism for CNgIc production (Kakes & Chardonnens, 2000; Olsen *et al.*, 2014). *T. occidentale* is not atypical among white clover's close congeners in this respect; several other *Trifolium* species also have been found to be polymorphic for CNgIc production while lacking the capacity to produce linamarase (e.g. *T. ambiguum*, *T. montanum*, *T. suffocatum*, *T. uniflorum*) (Gibson *et al.*, 1972; Olsen *et al.*, 2014). The most likely explanation for this pattern is that CNgIcs in these species serve functions in primary metabolism that are unrelated to cyanide release and herbivore deterrence. CNgIcs are produced in > 3000 plant species across > 130 families, and research over the last two decades has established important roles for them in physiological processes that do not involve cyanide release – most notably as storage and transport molecules for nitrogen (N) and glucose (Gleadow & Møller, 2014; Pičmanová *et al.*, 2015). For white clover evolution, it thus appears that the antiherbivore cyanogenic response evolved in part through co-option of CNgIc production for a new adaptive role as a chemical defense.

When functioning as storage and transport molecules, CNgIcs provide readily metabolized reserves of reduced-state N; these N reserves can be especially beneficial for plants growing in drought-prone or other stressful environments (Gleadow & Møller, 2014; Kooyers *et al.*, 2014). In the case of white clover, where climate-associated *Ac/ac* clines are well-documented throughout the species range (Daday, 1954, 1958; Kooyers & Olsen, 2012, 2013; Olsen *et al.*, 2013), both observational and experimental evidence suggest that some of this adaptive variation probably reflects selective pressures unrelated to cyanogenesis – particularly, plant fitness in drought-prone environments (Kooyers *et al.*, 2014). Thus, it seems likely that the presumed ancestral function of CNgIcs in primary metabolism was maintained in white clover even after they gained their well-documented adaptive function in herbivore deterrence.

Genome sequence comparisons between white clover and its diploid progenitors suggest that this is a very young polyploid species, having emerged around the time of the last major European glaciation *c.* 20 000 yr ago, when the progenitor species are expected to have had regions of range overlap in glacial refugia (Griffiths *et al.*, 2019). Consistent with this scenario, the white clover *Ac* locus shows very high sequence conservation with its

T. occidentale ortholog; the three CNglc synthesis genes and most intervening noncoding sequence show $\geq 97\%$ identity on average between species (Fig. 2). These similarity estimates would likely be even higher with a more complete version of the *T. occidentale* reference genome, as the current shotgun assembly has multiple regions of undetermined sequence within the *Ac* locus (see cross-hatched zones in Fig. 2).

Genes such as *Ac* that comprise metabolic clusters generally are believed to under strong selection to remain in close physical proximity within the genome to ensure inheritance of the complete biosynthetic pathway, thereby minimizing risk of accumulating toxic pathway intermediates (Takos & Rook, 2012; Gleadow & Møller, 2014). In a previous study of the white clover *Ac* cluster, we found that the three component genes were orthologous to and highly conserved with those of a CNglc cluster in the cyanogenic species *Lotus japonicus* (birdsfoot trefoil), which last shared an ancestor with *Trifolium c.* 50 Myr ago (Olsen & Small, 2018). Notably, on the timescale of that much older evolutionary divergence, there was essentially no conservation of the cluster outside the three highly conserved CNglc genes; gene order, transcript orientation and intervening sequences were all completely diverged. This pattern suggests that there is little overall selective constraint on noncoding portions of the CNglc metabolic cluster. In this light, the very high similarity we observe for the entire gene cluster between *T. repens* and *T. occidentale* (Fig. 2) further supports our conclusion that the white clover *Ac* locus is directly and recently descended from that progenitor species.

White clover is characterized by high genetic diversity despite its recent hybrid origin, and coalescent modeling has suggested that the species likely evolved through multiple independent *T. pallescens* \times *T. occidentale* hybridization events (Griffiths *et al.*, 2019). Given that *T. occidentale* carries the *Aclac* polymorphism, independent hybridizations involving *Ac*₋ and *acac* genotypes of that species could easily account for the origin of the white clover *Aclac* polymorphism. If true, this would mean that the gene-presentation and -absence alleles that characterize the well-studied *Aclac* polymorphism in white clover pre-date the origin of the species, making it a case of trans-specific adaptive polymorphism. However, molecular evolutionary analyses of the *Ac* locus and flanking sequences indicate that the genomic deletions that characterize *ac* alleles have evolved repeatedly, both in white clover and in its congeners that carry the *Aclac* polymorphism (Olsen *et al.*, 2013, 2014). Thus, genomic instability around the *Ac* locus may lend itself to repeated evolution of species-specific *ac* alleles even if the original white clover *Aclac* polymorphism was inherited directly from *T. occidentale*.

Linamarase and the *Li* locus

Although genetic mapping places the white clover *Li* locus unambiguously in the *T. pallescens* subgenome (linkage group 4P; Fig. 1; Table 1), the absence of the gene from the *T. pallescens* reference genome prevents sequence comparisons between the species. The gene's absence can most easily be explained in one of two ways. *T. pallescens* may carry the *Li/li* gene presence/absence

polymorphism, as occurs in white clover and other closely related species (e.g. *T. isthmocarpum*, *T. nigrescens*; Olsen *et al.*, 2014), with the reference genome representing a *lili* genotype. Alternatively, the gene may have been present at the time that *T. pallescens* hybridized with *T. occidentale* to create *T. repens*, but later lost in the *T. pallescens* lineage. Several factors favor the second explanation. First, neither linamarase nor the underlying *Li* gene have ever been identified in *T. pallescens* in surveys to date (Gibson *et al.*, 1972; Olsen *et al.*, 2014) (Table S1). Equally notably, CNglcs have not been detected in the species either, and linamarase has no known function other than in CNglc hydrolysis to release HCN; it thus would be implausible for a species to produce the CNglc-specific hydrolytic enzyme without CNglcs also being produced in at least some individuals. The restricted distribution of *T. pallescens* within cool alpine habitats further argues against the gene's present-day occurrence in the species. Cyanogenic *Trifolium* species (or cyanogenic populations of polymorphic species) occur primarily in warm temperate and hot Mediterranean climates (Olsen *et al.*, 2013, 2014); *T. pallescens* would thus be a unique exception in the genus if it were an alpine-adapted cyanogenic species. Based on all of these factors, the most likely scenario may be that *T. pallescens* was once cyanogenic (or polymorphic for cyanogenesis), and that this trait was lost during the species' post-Pleistocene range contraction into its present-day alpine environment.

It should be noted that a third potential explanation also exists for the occurrence of the *Li* gene in *T. repens* which we cannot exclude with our present data – namely, adaptive introgression from some other *Trifolium* species that carries this gene. Despite differences in ploidy between white clover and the two other *Trifolium* species that are known to produce linamarase (*T. nigrescens* and *T. isthmocarpum*, both $2n = 2x = 16$), each of these species can be hybridized with *T. repens* under controlled conditions, and crosses with *T. nigrescens* can produce fertile hybrids (Kazimierski & Kazimierska, 1972; Abberton, 2007). Thus, although it is not directly suggested by our genetic map data, an adaptive introgression scenario for the origin of *Li* in white clover also remains within the realm of possibility.

Allopolyploidy and cyanogenesis evolution in white clover

Speciation via allopolyploidy is widely recognized as a potential source of evolutionary novelty, with the combination of two or more parental species' genomes providing opportunities for dynamic remodeling of genome organization, gene content, gene expression networks and transposon activity (Soltis *et al.*, 2014). These changes provide a novel genomic toolkit that can potentially facilitate adaptation and ecological niche expansion (reviewed in Van de Peer *et al.*, 2017). White clover may well be an illustration of this phenomenon (Griffiths *et al.*, 2019), as the species occurs abundantly worldwide, in climates ranging from boreal to subtropical, despite its parental species' narrow ecological and geographical distributions. Although the role of cyanogenesis in white clover's ability to expand into such a wide environmental range is unknown, the fact that climate-associated cyanogenesis clines have evolved repeatedly throughout the

species range (reviewed by Kooyers & Olsen, 2012; Olsen *et al.*, 2013) suggests that selection on this trait plays an important role in the species' wide ecological amplitude. At the same time, it should be noted that cyanogenesis is unlikely to have been a novel trait that evolved uniquely in *Trifolium* with the origin of *T. repens*; as discussed above, several congeners also are cyanogenic. In this context, the dual-species origin of white clover cyanogenesis that we have documented in this study may best be viewed as one feature of the newly evolved species that, in combination with other emergent features of its allopolyploid toolkit, facilitated its global expansion and ecological success.

Acknowledgements

The authors thank Mike Dyer and other staff of the Washington University greenhouse facility for providing plant care; Andrew Griffiths and Roger Moraga (AgResearch NZ) for reference genome sequence information; Rachel Tan (AgResearch NZ) and Jennifer Barrett (Danforth Plant Science Center, St Louis) for guidance on genotyping-by-sequencing protocols; Marshall Wedger (Washington University) for assistance with QTL mapping scripts; and members of the Olsen lab group for helpful comments on the manuscript. Funding for the work was provided by NSF grants IOS-1557770 to KMO and DEB-1601641 to SJW, and by NSF graduate research fellowship DGE-1143954 to SJW. The authors declare no competing interests.

Author contributions

KMO conceived the project, analyzed sequence data, interpreted the results and wrote the manuscript; SJW created the mapping populations, generated GBS genotype data and contributed to writing; L-FL and DMG performed bioinformatics; DMG also constructed linkage maps and performed QTL mapping and subgenome data interpretation; and MLD, SRM and LLS performed cyanogenesis genotyping and phenotyping. All authors contributed comments on earlier drafts of the manuscript. DMG and SJW contributed equally to this work.

ORCID

David M. Goad  <https://orcid.org/0000-0001-8658-6660>
Kenneth M. Olsen  <https://orcid.org/0000-0002-8338-3638>

Data availability

Genotyping-by-sequencing reads have been submitted to NCBI SRA as aligned BAM files. The STL × GFL population is available under project ID PRJNA740324 (accessions SAMN19839204–SAMN19839729). The DMN × GFL population is under project ID PRJNA740343 (accessions SAMN19841524–SAMN19842038). Marker information and other genetic map data are provided in the online supplementary materials (Tables S8, S9).

References

- Abberton MT. 2007. Interspecific hybridization in the genus *Trifolium*. *Plant Breeding* 126: 337–342.
- Anssour S, Baldwin IT. 2010. Variation in antiherbivore defense responses in synthetic *Nicotiana* allopolyploids correlates with changes in uniparental patterns of gene expression. *Plant Physiology* 153: 1907–1918.
- Armstrong HE, Armstrong EF, Horton E. 1913. Herbage studies. II. Variation in *Lotus corniculatus* and *Trifolium repens* (Cyanophoric Plants). *Proceedings of the Royal Society of London. Series B: Biological Sciences* 86: 262–269.
- Badr A, Sayed-Ahmed H, El-Shanshoury A, Watson IE. 2002. Ancestors of white clover (*Trifolium repens* L.), as revealed by isozyme polymorphisms. *Theoretical and Applied Genetics* 106: 143–148.
- Barrett B, Griffiths A, Schreiber M, Ellison N, Mercer C, Bouton J, Ong B, Forster J, Sawbridge T, Spangenberg G *et al.* 2004. A microsatellite map of white clover. *Theoretical and Applied Genetics* 109: 596–608.
- Bilton TP, Schofield MR, Black MA, Chagné D, Wilcox PL, Dodds KG. 2018. Accounting for errors in low coverage high-throughput sequencing data when constructing genetic maps using biparental outcrossed populations. *Genetics* 209: 65–76.
- Brachi B, Meyer CG, Villoutreix R, Platt A, Morton TC, Roux F, Bergelson J. 2015. Coselected genes determine adaptive variation in herbivore resistance throughout the native range of *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* 112: 4032–4037.
- Briggs D, Walters SM. 2016. *Plant variation and evolution*. Cambridge, UK: Cambridge University Press. doi: <https://doi.org/10.1017/CBO9781139060196>.
- Broman KW, Wu H, Sen S, Churchill GA. 2003. *R/qtl*: QTL mapping in experimental crosses. *Bioinformatics* 19: 889–890.
- Casey NM, Milbourne D, Barth S, Febrer M, Jenkins G, Abberton MT, Jones C, Thorogood D. 2010. The genetic location of the self-incompatibility locus in white clover (*Trifolium repens* L.). *Theoretical and Applied Genetics* 121: 567–576.
- Daday H. 1954. Gene frequencies in wild populations of *Trifolium repens* L. II. Distribution by altitude. *Heredity* 8: 377–384.
- Daday H. 1958. Gene frequencies in wild populations of *Trifolium repens* L. III. World distribution. *Heredity* 12: 169–184.
- Edger PP, Heidel-Fischer HM, Bekeert M, Rota J, Glöckner G, Platts AE, Heckel DG, Der JP, Wafula EK, Tang M *et al.* 2015. The butterfly plant arms-race escalated by gene and genome duplications. *Proceedings of the National Academy of Sciences, USA* 112: 8362–8366.
- Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, Mitchell SE. 2011. A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PLoS ONE* 6: e19379.
- Feigl F, Anger V. 1966. Replacement of benzidine by copper ethylacetoacetate and tetra base as spot-test reagent for hydrogen cyanide and cyanogen. *The Analyst* 91: 282–284.
- Futuyma DJ, Kirkpatrick M. 2017. *Evolution, 4th edn*. Sunderland, MA, USA: Sinauer. [WWW document] URL <https://evolution4e.sinauer.com/> [accessed 8 March 2021].
- Gibson PB, Barnett OW, Gillingham JT. 1972. Cyanoglucoside and hydrolyzing enzyme in species related to *Trifolium repens*. *Crop Science* 12: 708–709.
- Gleadow RM, Møller BL. 2014. Cyanogenic glycosides: synthesis, physiology, and phenotypic plasticity. *Annual Review of Plant Biology* 65: 155–185.
- Griffiths AG, Barrett BA, Simon D, Khan AK, Bickerstaff P, Anderson CB, Franzmayr BK, Hancock KR, Jones CS. 2013. An integrated genetic linkage map for white clover (*Trifolium repens* L.) with alignment to *Medicago*. *BMC Genomics* 14: 388.
- Griffiths AG, Moraga R, Tausen M, Gupta V, Bilton TP, Campbell MA, Ashby R, Nagy I, Khan A, Larking A *et al.* 2019. Breaking free: the genomics of allopolyploidy-facilitated niche expansion in white clover. *Plant Cell* 31: 1466–1487.
- Haley CS, Knott SA. 1992. A simple regression method for mapping quantitative trait loci in line crosses using flanking markers. *Heredity* 69: 315–324.
- Huang P, Feldman M, Schroder S, Bahri BA, Diao X, Zhi H, Estep M, Baxter I, Devos KM, Kellogg EA. 2014. Population genetics of *Setaria viridis*, a new model system. *Molecular Ecology* 23: 4912–4925.

- Hughes MA. 1991. The cyanogenic polymorphism in *Trifolium repens* L. (white clover). *Heredity* 66: 105–115.
- Kakes P. 1985. Linamarase and other β -glucosidases are present in the cell walls of *Trifolium repens* L. leaves. *Planta* 166: 156–160.
- Kakes P, Chardonnens AN. 2000. Cyanotypic frequencies in adjacent and mixed populations of *Trifolium occidentale* Coombe and *Trifolium repens* L. are regulated by different mechanisms. *Biochemical Systematics and Ecology* 28: 633–649.
- Kakes P, Hakvoort HWJ. 1994. On the origin of the cyanogenic polymorphism in *Trifolium repens* L. *Journal of Evolutionary Biology* 7: 201–215.
- Kazimierski T, Kazimierska EM. 1972. Badania mieszańców w rodzaju *Trifolium* L. IV. Cytogenetyka mieszańca *Trifolium repens* L. \times *T. isthmocarpum* Brot. [Investigations on hybrids of the genus *Trifolium* L. IV. Cytogenetics of the cross *T. repens* L. \times *T. isthmocarpum* Brot.]. *Acta Societatis Botanicorum Poloniae* 41: 129–147.
- Keith RA, Mitchell-Olds T. 2019. Antagonistic selection and pleiotropy constrain the evolution of plant chemical defenses. *Evolution* 73: 947–960.
- Kooyers N, Bakken B, Ungerer M, Olsen K. 2018. Freeze-induced cyanide toxicity does not maintain the cyanogenesis polymorphism in white clover (*Trifolium repens*). *American Journal of Botany* 105: 1224–1231.
- Kooyers NJ, Gage LR, Al-Lozi A, Olsen KM. 2014. Aridity shapes cyanogenesis cline evolution in white clover (*Trifolium repens* L.). *Molecular Ecology* 23: 1053–1070.
- Kooyers NJ, Olsen KM. 2012. Rapid evolution of an adaptive cyanogenesis cline in introduced North American white clover (*Trifolium repens* L.). *Molecular Ecology* 21: 2455–2468.
- Kooyers NJ, Olsen KM. 2013. Searching for the bull's eye: agents and targets of selection vary among geographically disparate cyanogenesis clines in white clover (*Trifolium repens* L.). *Heredity* 111: 495–504.
- Lai D, Maimann AB, Macea E, Ocampo CH, Cardona G, Pičmanová M, Darbani B, Olsen CE, Debouck D, Raatz B *et al.* 2020. Biosynthesis of cyanogenic glucosides in *Phaseolus lunatus* and the evolution of oxime-based defenses. *Plant Direct* 4: e00244.
- Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25: 1754–1760.
- Lowry DB, Popovic D, Brennan DJ, Holeski LM. 2019. Mechanisms of a locally adaptive shift in allocation among growth, reproduction, and herbivore resistance in *Mimulus guttatus*. *Evolution* 73: 1168–1181.
- Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet* 17: 10–12.
- Mithöfer A, Boland W. 2012. Plant defense against herbivores: chemical aspects. *Annual Review of Plant Biology* 63: 431–450.
- Morant AV, Jørgensen K, Jørgensen C, Paquette SM, Sánchez-Pérez R, Møller BL, Bak S. 2008. beta-Glucosidases as detonators of plant chemical defense. *Phytochemistry* 69: 1795–1813.
- Olsen KM, Hsu S-C, Small LL. 2008. Evidence on the molecular basis of the *Ac/ac* adaptive cyanogenesis polymorphism in white clover (*Trifolium repens* L.). *Genetics* 179: 517–526.
- Olsen KM, Kooyers NJ, Small LL. 2013. Recurrent gene deletions and the evolution of adaptive cyanogenesis polymorphisms in white clover (*Trifolium repens* L.). *Molecular Ecology* 22: 724–738.
- Olsen KM, Kooyers NJ, Small LL. 2014. Adaptive gains through repeated gene loss: parallel evolution of cyanogenesis polymorphisms in the genus *Trifolium* (Fabaceae). *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 369: 20130347.
- Olsen KM, Small LL. 2018. Micro- and macroevolutionary adaptation through repeated loss of a complete metabolic pathway. *New Phytologist* 219: 757–766.
- Olsen KM, Sutherland BL, Small LL. 2007. Molecular evolution of the *Lilli* chemical defence polymorphism in white clover (*Trifolium repens* L.). *Molecular Ecology* 16: 4180–4193.
- Pichersky E, Lewinsohn E. 2011. Convergent evolution in plant specialized metabolism. *Annual Review of Plant Biology* 62: 549–566.
- Pičmanová M, Neilson EH, Motawia MS, Olsen CE, Agerbirk N, Gray CJ, Flitsch S, Meier S, Silvestro D, Jørgensen K *et al.* 2015. A recycling pathway for cyanogenic glycosides evidenced by the comparative metabolic profiling in three cyanogenic plant species. *Biochemical Journal* 469: 375–389.
- Rimmer A, Phan H, Mathieson I, Iqbal Z, Twigg SRF, Consortium W, Wilkie AOM, Mcvean G, Lunter G. 2014. Integrating mapping-, assembly- and haplotype-based approaches for calling variants in clinical sequencing applications. *Nature Genetics* 46: 912–918.
- Santangelo JS, Thompson KA, Cohan B, Syed J, Ness RW, Johnson MTJ. 2020. Predicting the strength of urban-rural clines in a Mendelian polymorphism along a latitudinal gradient. *Evolution Letters* 4: 212–225.
- Soltis PS, Liu X, Marchant DB, Visger CJ, Soltis DE. 2014. Polyploidy and novelty: Gottlieb's legacy. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 369: 20130351.
- Takos AM, Rook F. 2012. Why biosynthetic genes for chemical defense compounds cluster. *Trends in Plant Science* 17: 383–388.
- Thompson KA, Renaudin M, Johnson MTJ. 2016. Urbanization drives the evolution of parallel clines in plant populations. *Proceedings of the Royal Society of London. Series B: Biological Sciences* 283: 20162180.
- Torkamaneh D, Laroche J, Bastien M, Abed A, Belzile F. 2017. Fast-GBS: a new pipeline for the efficient and highly accurate calling of SNPs from genotyping-by-sequencing data. *BMC Bioinformatics* 18: 1–7.
- Van de Peer Y, Mizrahi E, Marchal K. 2017. The evolutionary significance of polyploidy. *Nature Reviews. Genetics* 8: 206–216.
- Wager A, Li X. 2018. Exploiting natural variation for accelerating discoveries in plant specialized metabolism. *Phytochemistry Reviews* 17: 17–36.
- Whitlock R, Hipperson H, Mannarelli M, Burke T. 2008. A high-throughput protocol for extracting high-purity genomic DNA from plants and animals. *Molecular Ecology Resources* 8: 736–741.
- Williams WM, Ellison NW, Ansari HA, Verry IM, Hussain SW. 2012. Experimental evidence for the ancestry of allotetraploid *Trifolium repens* and creation of synthetic forms with value for plant breeding. *BMC Plant Biology* 12: 55.
- Xu S, Brockmüller T, Navarro-Quezada A, Kuhl H, Gase K, Ling Z, Zhou W, Kreitzer C, Stanke M, Tang H *et al.* 2017. Wild tobacco genomes reveal the evolution of nicotine biosynthesis. *Proceedings of the National Academy of Sciences, USA* 114: 6133–6138.
- Zhuang X, Köllner TG, Zhao N, Li G, Jiang Y, Zhu L, Ma J, Degenhardt J, Chen F. 2012. Dynamic evolution of herbivore-induced sesquiterpene biosynthesis in sorghum and related grass crops. *The Plant Journal* 69: 70–80.

Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

Fig. S1 Synteny plots comparing marker order in the white clover linkage maps to the locations of their best BLAST hit in the *Medicago truncatula* genome, for the DG population and the SG population.

Fig. S2 QTL peak for *Ac/ac* in the DG mapping population.

Fig. S3 QTL peak for *Ac/ac* in the SG mapping population.

Fig. S4 QTL peak for *Lilli* in the DG mapping population.

Fig. S5 QTL peak for *Lilli* in the SG mapping population.

Fig. S6 QTL peaks for cyanogenesis in the DG mapping population.

Fig. S7 QTL peak for cyanogenesis in the SG mapping population.

Table S1 *Ac* and *Li* phenotypes and genotypes for *Trifolium pallelescens* accessions.

Table S2 Percentage sequence identity between a sample of *Trifolium occidentale* and *Trifolium pallescens* orthologs.

Table S3 Parentage and cyanogenesis genotyping for the DG mapping population.

Table S4 Parentage and cyanogenesis genotyping for the SG mapping population.

Table S5 Primer combinations for *Ac* and *Li* genotyping, and GBS PCR primers for final library amplification and Illumina sequencing.

Table S6 White clover GBS common and barcode adapters.

Table S7 Number of markers on each linkage group with best BLAST hit to either the *Trifolium occidentale* or *Trifolium pallescens* reference genomes.

Table S8 GBS marker data for the DG population.

Table S9 GBS marker data for the SG population.

Table S10 GenBank accessions of best BLAST hits for white clover cyanogenesis loci in the *Trifolium occidentale*, *Trifolium pallescens* and *Trifolium repens* draft genomes.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the New Phytologist Central Office.



About New Phytologist

- *New Phytologist* is an electronic (online-only) journal owned by the New Phytologist Foundation, a **not-for-profit organization** dedicated to the promotion of plant science, facilitating projects from symposia to free access for our Tansley reviews and Tansley insights.
- Regular papers, Letters, Viewpoints, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as ready' via *Early View* – our average time to decision is <26 days. There are **no page or colour charges** and a PDF version will be provided for each article.
- The journal is available online at Wiley Online Library. Visit **www.newphytologist.com** to search the articles and register for table of contents email alerts.
- If you have any questions, do get in touch with Central Office (np-centraloffice@lancaster.ac.uk) or, if it is more convenient, our USA Office (np-usaoffice@lancaster.ac.uk)
- For submission instructions, subscription and all the latest information visit **www.newphytologist.com**